

Adam Para,
Fermilab, December 10, 2009
Non-Standard Neutrino Interactions Workshop
Madrid

WWW.MINSIS

WHAT

WHY

WHERE

WHEN

WHO

Neutrino Physics circa 1995

- Tantalizing suggestions:
 - Neutrinos have mass?
 - Flavor non-conservation?
 - Neutrino oscillations?
- Two prongs:
 - Natural: large mass difference, very small mixing angle
↔ Short baseline oscillation experiments
 - Unnatural: large mixing angle, very small mass difference
↔ Long baseline oscillation experiments

Short baseline experiments, 1995

- CHORUS/NOMAD: ν_τ appearance probability $< 10^{-4}$
- Mixing angles are expected (naturally) to be very small, hence
- Second generation of short baseline oscillation experiments: huge amount of detailed simulations and detector design work:
 - COSMOS
 - TENOR
 - TOSCA
 - NAUSICAA
 - ESTAR

SuperKamiokande bombshell

- Neutrino oscillations
- Very small mass difference
- Neutrino oscillations very strongly suppressed (~absent) at short baseline
- Sudden death of all short baseline oscillation experiments

Particle Physics circa 2010

- Neutrino oscillations well established, mass differences and mixing angles pretty well known
- Standard model expected to be incomplete, new physics expected in the range \sim few hundred GeV
- Standard model spectacularly successful: no detectable deviation (other than the neutrino masses). What does it mean?
 - Mass/energy scale of the new physics much larger? Multi TeV or higher?
 - Some symmetries/cancellations reducing the contribution of new physics to the investigated processes?


Broader Search for the New Physics? Neutrinos?

- (Almost?) every extension of the standard model leads to detectable effects in the neutrino sector
- Neutrino processes involving the third generation (taus/tau neutrinos) may be particularly complementary to the other searches for the non-standard interactions
- Neutrino oscillations may hide the effects of the new interactions (Nobel Prize of yesterday a tomorrow's background)
- Remember the fate of the short baseline oscillation experiments?? Neutrino oscillations are 'not observable' at the short distance: → particularly sensitive probe of new physics

Search for $\nu_\mu(\nu_e) \leftrightarrow \nu_\tau$ Oscillations with a Detector Based on a Emulsion-Silicon Target

J.J. Gomez-Cadenas^{1,2} and J.A. Hernando^{2,3}

It is possible to improve the limit on the ν_μ to ν_τ conversion by about two orders of magnitude, or discover the new physics (neutrino oscillations) using the state of the art experimental techniques and planned NuMI neutrino beam (1995)



From 1995 to 2010

- Tau neutrino interactions observed (DONUT)
- New emulsion techniques proven to be superior, cheaper and easier to the bulk emulsions (DONUT)
- Dramatic increase in the speed and efficiency of the automatic scanning techniques
- Huge, 2 kton, tau appearance experiment constructed and operating (OPERA)
- Silicon tracking (strips and/or pixels) well established technology. Huge area silicon detectors constructed and operating (CDF, D0, CMS, ATLAS)
- High intensity neutrino beam (NuMI) constructed and operating. Major upgrade underway, to be completed in ~ 3 years

MINSIS: Main Injector Non Standard Neutrino Interactions Search?



- Proposition:
 - use the upgraded NuMI neutrino beam
 - take advantage of the huge investment in the design of the short baseline neutrino oscillation
 - Take advantage the enormous progress of the experimental techniques (emulsions, scanning, silicon)
 - Take advantage of the suppression of the oscillation background at short distances

to extend the search for the rare ν_{μ} -to- ν_{τ} conversion with the sensitivity up to $\sim 10^{-6}$

MINSIS: three fundamental (and related) questions:

- Is it possible (rates, efficiencies, backgrounds) ?
Beam (intensity, spectrum, composition), detailed detector design, analysis techniques..
- Is it affordable ?
- Is it important/interesting enough to bother ?



NuMI Beam and Near Detector Hall



NuMI Neutrino Beam

Variable Energy Neutrino Beam

Low Energy Beam



Pions with
 $p_T=300$ MeV/c and
 $p=5$ GeV/c
 $p=10$ GeV/c
 $p=20$ GeV/c

Vary ν beam energy by
sliding the target in/out
of the 1st horn

High Energy Beam





figure courtesy Ž. Pavlović

Horn 2


NuMI Neutrino Beam

- Constructed and operating (MINOS, MINERvA)
- Flexible design, adjustable beam energy and spectrum
- 370 kW power
- Major intensity upgrade for NOvA (700 kW)
- Fix the design, limit the flexibility of the future beam, given the well defined physics program (NOvA)
- → If some new experiment is to be contemplated it is important to include the possible requirements into the design process for the upgraded NuMI beam (for example: energy spectrum, antineutrino component)

The MINOS Experimental Hall, as built



MINOS Near Detector Hall, Now



MINERvA

Near Term Running Plan

Draft 2010-13 Fermilab Accelerator Experiments' Run Schedule

Typically Revised Annually - This Version from October, 2009

Calendar Year		2010	2011	2012	2013
Tevatron Collider		CDF & DZero	CDF & DZero	OPEN	OPEN
Neutrino Program	B	MiniBooNE	MiniBooNE		OPEN
		OPEN	OPEN		MicroBooNE
	MI	MINOS	MINOS		OPEN
		MINERvA	MINERvA		MINERvA
		ArgoNeuT			
				NOvA	NOvA
SY 120	MT	Test Beam	Test Beam		Test Beam
	MC	OPEN	OPEN		OPEN
	NM4	E-906/Drell-Yan	E-906/Drell-Yan		E-906/Drell-Yan

NoVA expected to run from 2013 till 2020 or so. MINERvA long term future less well defined. But MINSIS can be built and run concurrently with the both experiments.

Achieving the Sensitivity: Backgrounds

- The dominant beam related backgrounds:
 - Tau neutrinos component of the neutrino beam

Tau neutrinos are produced from decays of charm particles produced by the primary protons.

$\sigma_{\text{charm}} \sim 30 \mu\text{b}$ at 450 GeV, $\sim 5 \mu\text{b}$ at 120 GeV;
 $\nu_{\tau}/\nu_{\mu} < 10^{-7}$

 - Anti-charm produced by antineutrinos in the beam

May require a reduction of the intrinsic antineutrino component of the beam. Plug?

Achieving the Sensitivity: Detection Efficiency, Detector Backgrounds

- Requires detailed detector optimization, including the beam design
- A lot of practical experience from OPERA design, construction and operation. Need good feedback

	CHORUS	OPERA	MINSIS
Target Mass	800kg	1300ton	10 ton?
Emulsion Mass	800kg	30ton	1~10 ton
Cost		~100 M\$????

Emulsion Scanning Technology

	CHORUS	OPERA	MINSIS
Scan Area	1 m ² /4year	100m ² /year	20m ² ???
# of events	500K/4year	4K/year	10M
# of Films	600	9.3M	100K~10K
System	NTS / UTS	SUTS	SUTS
Speed	~1cm ² /h	100cm ² /h	100cm ² /h
LOAD	5 Years	1 Year	3 months

Much faster system ~5000cm²/h (SQTS) for dark matter, double beta experiment and Muon radiography under development in Nagoya. (M. Nakamura)

Analysis - Detector Design Interactions

- Past mindset: Scanning is a bottleneck, need a detector capable of finding the interactions and directing the measurement process
- Future mindset?: self contained emulsion detector, complete scan/analysis of the entire volume? Huge statistics of neutrino events as a by-product with simpler detectors ?

By-products?

- Charm Physics??

D0-D0bar Oscillation?

Charmed penta-quark study ?

Charmed hadron mean free path ?

Charmed Nucleus study?

Initial thoughts of M. Nakamura. Very important aspect of the experiment.

Thoughts on the Detector Design, Optimization, Protoyping...

- Start with the OPERA design
- Evaluate the background rejection capability
- Complement with silicon tracker (a la ESTAR)
- If necessary make the lead plates much thinner (50 microns steel?)
- Construct prototypes, use the existing NuMI beam for evaluation
-

Very Preliminary Impressions

- A new experiment searching for the tau appearance at short baseline using the NuMI neutrino beam with the sensitivity of the order of 10^{-6} is possible and quite realistic.
- Detailed and very careful studies necessary to optimize the ultimate experiment and to determine its physics potential.
- The present OPERA experiment the primary source of inspiration and of the critical evaluation of the detection efficiency and background rejection estimates.
- But.. Construction, operation and analysis of such an experiment does require a significant effort

Is It Worth Doing?? Part I

- A positive result would have a huge impact on our understanding of the particle physics. It would be a proof of some physics beyond the current standard model. It would be even better if some specific scenarios could be established? Is it a consequence of some sterile neutrinos? Or leptoquarks? Or SUSY? Or charged higgses? Is this possible to tell?? How does it complement the possible discoveries at the LHC? Or, perhaps, on the contrary, taking all of the existing limits and some sensible assumptions, the existence of such a process at the level of 10^{-6} or higher can be already excluded?

Is It Worth Doing?? Part II

- Suppose that after a heroic effort the experiment will demonstrate that the tau appearance process is suppressed by more than a factor of 10^{-6} with respect to standard neutrino interactions. Will anybody care? What sort of models will it exclude? What phase space in the parameters space? How likely is it that such a limit will be interesting at the time it can be established (say 2017?)

Establishing of the physics motivation for such a proposal is the most important step towards such a putative experiment.

Synergies

- 'OPERA crowd' - large group, expertise, existing detector, large investment in emulsion scanning and analysis, large amount of emulsion unused after the end of the experiment.. MINSIS = ActII of OPERA?
- 'Short baseline oscillation search crowd' - ESTAR, COSMOS, TOSCA people. Efficiency and background rejection studies of direct applicability to MINSIS, possible enhancement of the OPERA-like detector technology
- 'Neutrino factory near detector crowd' - MINSIS as an R&D/prototype project for the future experiment at the neutrino factory